

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# AMBITIOUS™

## PLANT GROWTH REGULATOR

**ACTIVE CONSTITUENT: 10 g/L FORCHLORFENURON**

Plant growth regulator for increasing berry size in Table Grapes, and fruit size in Apples and Cherries

**IMPORTANT: Read attached leaflet before use.**



NET CONTENTS

**1L, 5L**

**DISTRIBUTED BY:** 7 Worlds Ag – Grochem Australia Pty Ltd

Suite 1, Level 3, 262 Lorimer Street,  
Port Melbourne VIC 3207  
Telephone 1800 777 068  
APVMA Approval No. 65381/142777



**7 WORLDS AG**



**7 WORLDS AG**

7worlds.com.au | for all enquiries 1800 777 068

CONTENTS **1L, 5L**

## **DIRECTIONS FOR USE**

### **RESTRAINTS**

For PROFESSIONAL use only.

DO NOT apply by air.

DO NOT spray if rain is expected within 2 hours.

DO NOT allow bystanders to come into contact with the spray cloud.

### **SPRAY DRIFT RESTRAINTS**

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift).

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

<b>CROP</b>	<b>Rate per 100 L water</b>	<b>CRITICAL COMMENTS</b>
<b>Table Grapes</b> Thompson Seedless Menindee Seedless Red Globe	50-100mL	Apply at 4 to 6mm berry size.
<b>Apples</b> Royal Gala, Fuji	100 mL	Apply when king fruit size is 10mm. DO NOT apply more than one application per season. DO NOT apply later than crop growth stage BBCH 71.
Cherries	200 – 300 mL	Apply when fruit are at green fruit stage (7-10 days after petal or shuck fall) or at straw fruit colour. DO NOT apply more than one application per season. DO NOT apply later than crop growth stage BBCH 85.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## **WITHHOLDING PERIODS**

### **HARVEST:**

**Apples, Cherries, Grapes: NOT REQUIRED WHEN USED AS DIRECTED**

### **GRAZING:**

**DO NOT GRAZE TREATED VINEYARDS OR ORCHARDS OR CUT TREATED AREAS FOR STOCK FEED UNTIL AFTER COMPLETION OF HARVEST**

**EXPORT HARVEST INTERVAL (EHI): NOT REQUIRED WHEN USED AS DIRECTED**

## **GENERAL INSTRUCTIONS**

- Ambitious Plant Growth Regulator is a very active plant growth regulator with cytokinin activity which accelerates cell division, cell number and cell size. Promotes fruit enlargement as a result.
- The product is not translocated in plant tissue (e.g. from treated leaves to fruit), it is sufficient to spray only the fruit by direct application.
- Direct spray to the bunches and fruit. Bunches or fruit must be thoroughly wetted. The formation of drip points must be avoided.
- Run off and drip points may result in marks or rings on berry skin or lop sided, misshapen apples.
- Avoid application in midday heat.

### Grapes

- Apply after post-bloom shatter period to avoid increased fruit set.
- Gibberellic Acid – Good berry thinning is essential when using Ambitious Plant Growth Regulator. Best results are obtained when using Ambitious in addition to a GA3 program.
- On established vines apply in water rates between 500-1000L per ha. Spray volumes will depend on crop load and spray equipment. Reduce water volumes on young vines.
- Higher rates of Ambitious Plant Growth Regulator may change berry shape to a more spherical form.
- Ambitious Plant Growth Regulator can be applied:
  - o Directed spray to young fruit or bunches, with spray volumes in the range of 500 L/ha (single fruit treatment). Spray volumes depend on crop load, spray equipment used, nozzle type and spray pressure. Bunches have to be thoroughly wetted. Formation of drip points must be avoided.
  - o Overall spray on the whole vines by means of a mist blower, etc. (e.g. air blast spraying), with spray volumes of around 1000 L/ha depending on vine height, spacing of vines and equipment used (nozzle type and size etc.).
- Apply as a fine mist. Complete coverage of the fruit is essential, and all bunches have to be wetted thoroughly. Apply at a volume of between 500- 1000 L water per hectare on established vines. Use proportionally less on younger vines. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- The required spray volume will change, and the sprayer set up and operation may also need to be changed as the crop grows.
- Avoid run-off and formation of drip-points, which may lead to unwanted marks/rings on the berry skin. Apply preferably during the morning or late afternoon hours. Avoid the midday heat. Avoid spraying if rain is expected. As Ambitious Plant Growth Regulator is not systemic, and applications are made some weeks before new bud formation occurs. Ambitious Plant Growth Regulator cannot directly influence crop load in the following year. However, with the use of Ambitious Plant Growth Regulator, the productive requirements of the treated plants are increased. For this reason, it is important to adjust fruit load, nutritional state and water balance of the plants accordingly.
- Apply Ambitious Plant Growth Regulator after termination of post bloom shatter period as otherwise Ambitious Plant Growth Regulator may increase fruit set.

#### Apples and Cherries

- To increase fruit size of small fruit varieties.
- Application will not compensate for other limiting factors, such light crop load, drought, heat, nutritional, disease or other stress on crop.
- Apply as a fine, dilute spray in sufficient water to provide good coverage for the target canopy at crop stage, include a non-ionic surfactant such as Pomade™. Avoiding sprayer set ups and water rates that result in drip points.
- Apply at a target fruit size of 10mm king fruit.
- Apply under warm conditions, with high for the day 200C or greater. Avoid applying in the heat of the day.
- Product does not translocate across fruit. Fruit ridging or deformations are possible, care should be taken to ensure even coverage and do not use above label rate.
- Fruit color development may be delayed in Fuji.
- Do not apply when rain is expected.

#### Mixing

Thoroughly clean out spray container with clean water. Use correct measuring device and do not mix more than is needed. To prevent foaming and ensure adequate mixing, add part of water to the spray tank, then product, mix and add reminding water. Agitate whilst mixing and applying.

#### Compatibility

Ambitious Plant Growth Regulator is not compatible with other insecticides, fungicides, herbicides or other than specified adjuvants. Ambitious Plant Growth Regulator may be used in a tank mix with Gibberellic Acid.

#### Wetting Agent

A non-ionic wetting agent such as Pomade™ or Spray Aid may be added to the tank.

#### RESISTANCE WARNING

##### INTEGRATED PEST MANAGEMENT

Cherries: Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas.

#### PRECAUTIONS

DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

#### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plant/crops, cropping lands or pastures.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life.

DO NOT contaminate streams, rivers, watercourses with the chemical or used containers.

#### STORAGE AND DISPOSAL

FLAMMABLE – keep away from flames or heat.

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a cool (below 20°C), dry place out of reach of children. Do not store in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site.

If recycling, replace cap and return clean containers to recycler or designated collection point.  
If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**SAFETY DIRECTIONS**

Will irritate the eyes and skin. Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. When using together with other products, consult their label safety directions. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766.

**SAFETY DATA SHEET**

For additional information, refer to SDS which is available from the supplier.

**CONDITIONS OF SALE**

This high-quality product is suitable for the purpose advised on the label but as Gro-Chem cannot exercise control over mixing or applications, all conditions and warranties, as to its use for any purpose, are excluded and no responsibility will be accepted by Gro-Chem for any damage or injury whatsoever arising from its storage or application.

APVMA Approval No.: 65381/142777

**Highly flammable liquid and vapour. Causes serious eye irritation. Suspected of causing cancer. Harmful to aquatic life with long lasting effects.**

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Keep away from heat, sparks, open flames and hot surfaces. - No smoking. Ground/bond container and receiving equipment. Use explosion-proof electrical equipment. Use only non-sparking tools. Take precautionary measures against static discharge. Avoid breathing fumes, mists, vapours or spray. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Wear protective gloves, protective clothing and eye or face protection.

**IN A TRANSPORT EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**

**For specialist advice in an emergency only, call  
CHEMTREC: 1800 862 115  
ALL HOURS – AUSTRALIA**



**7 WORLDS AG**

**DISTRIBUTED BY:**

**7 Worlds Ag – Grochem Australia Pty Ltd**

Suite 1, Level 3, 262 Lorimer Street

Port Melbourne VIC 3207

Phone: 1800 777 068