

Grochem Dodine is group U12 fungicide that has both curative and protective properties.

As a cell membrane disruptor, it has excellent curative protective properties against Black spot (Scab) in both Apples and Pears, when applied within 36 hours after the start of an infection period, it penetrates the leaf surface to eradicate the black spot or scab infection. Grochem Dodine also has superior protective properties against Peach Leaf Curl and Blossom Blight.



Apple and Pears (Black Spot / Scab)

PROTECTIVE PROGRAMME: Following a copper spray at green tip, apply at pink bud (apples) or white bud (pears) followed by 2 cover sprays 10 days apart, a further 2 cover sprays 10-14 days apart and a fifth cover spray after a further 21 days and then at 4-6 week intervals depending on weather and presence of infection.

CURATIVE PROGRAMME: Apply within 36 hours of the start of an infection period (eg after rain) which is favourable to development of black spot.

Peaches and Nectarines (Peach Leaf Curl and Blossom Blight)

Apply at bud swell, early bloom and petal fall. DO NOT apply after petal fall.

INSTRUCTIONS FOR DETAILS

RATE: In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.				CRITICAL COMMENTS: For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
CROP	DISEASE	RATE/100L	WITHOLDING	CRITICAL COMMENTS
Apples	Black Spot (Venturia inaequalis)	80ml	5 days	<p>PROTECTIVE PROGRAMME: Following a copper spray at green tip, apply at pink bud (apples) or white bud (pears) followed by 2 cover sprays 10 days apart, a further 2 cover sprays 10-14 days apart and a fifth cover spray after a further 21 days and then at 4-6 week intervals depending on weather and presence of infection.</p> <p>CURATIVE PROGRAMME: Apply within 36 hours of the start of an infection period (eg after rain) which is favourable to development of black spot. This fungicide is subject to resistance strategy. See General Instructions for details.</p>
Pears	Black Spot (Venturia pirina)	120ml		
Peaches	Peach Leaf Curl (Taphrina deformans)	100ml	-	Apply at bud swell, early bloom and petal fall. DO NOT apply after petal fall.
Nectarines	Blossom Blight (Monilinia spp)	200ml		

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

For more information, contact your local 7 Worlds Ag representative

www.7worlds.com.au

for all enquiries 1800 777 068

DODINE